

Author's Accepted Manuscript

Hunting and Hallucinogens: The use psychoactive and other Plants to improve the Hunting ability of dogs

Bradley C. Bennett, Rocío Alarcón



www.elsevier.com/locate/jep

PII: S0378-8741(15)00367-0
DOI: <http://dx.doi.org/10.1016/j.jep.2015.05.035>
Reference: JEP9537

To appear in: *Journal of Ethnopharmacology*

Received date: 27 February 2015
Revised date: 20 May 2015
Accepted date: 20 May 2015

Cite this article as: Bradley C. Bennett, Rocío Alarcón, Hunting and Hallucinogens: The use psychoactive and other Plants to improve the Hunting ability of dogs, *Journal of Ethnopharmacology*, <http://dx.doi.org/10.1016/j.jep.2015.05.035>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting galley proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Hunting and Hallucinogens: The Use Psychoactive and Other Plants to Improve the Hunting Ability of Dogs

Bradley C. Bennett^a and Rocío Alarcón^b

^aDepartment of Biological Sciences, Florida International University, Miami, FL 33199 USA

(bennett@fiu.edu, 305-348-3586)

^bIamoe Centre. Via El Pindo, Coca Ecuador (rocioalarcon73@gmail.com)

Abstract

Ethnopharmacological relevance

Cultures throughout the world give plants to their dogs in order to improve hunting success. These practices are best developed in lowland Ecuador and Peru. There is no experimental evidence for the efficacy of these practices nor critical reviews that consider possible pharmacological effects on dogs based on the chemistry of the ethnovertebrary plants.

Aim

This review has three specific aims: 1. Determine what plants the Ecuadorian Shuar and Quichua give to dogs to improve their hunting abilities, 2. Determine what plants other cultures give to dogs for the same purpose, and 3. Assess the possible pharmacological basis for the use of these plants, particularly the psychoactive ones?

Methods

We gathered Shuar (Province of Morona-Santaigo) and Quichua (Napo and Orellano Provinces) data from our previous publications and field notes. All specimens were vouchered and

deposited in QCNE with duplicates sent to NY and MO. Data presented from other cultures derived from published studies on ethnoveterinary medicine. Species names were updated, when necessary, and family assignments follow APG III (Angiosperm Phylogeny Group 2009). Chemical data were found using PubMed and SciFinder.

Results

The Shuar and Quichua of Ecuador use at least 22 species for ethnoveterinary purposes, including all but one of their principal hallucinogens. Literature surveys identified 43 species used in other cultures to improve hunting ability. No published studies have examined the pharmacological active of these plant species in dogs. We, thus, combined phytochemical data with the ethnobotanical reports of each plant and then classified each species into a likely pharmacological category: depuratives/deodorant, olfactory sensitizer, ophthalmic, or psychoactive.

Conclusions

The use of psychoactive substances to improve a dog's hunting ability seems counterintuitive, yet its prevalence suggests that it is both adaptive and that it has an underlying pharmacological explanation. We hypothesize that hallucinogenic plants alter perception in hunting dogs by diminishing extraneous signals and by enhancing sensory perception (most likely olfaction) that is directly involved in the detection and capture of game. If this is true, plant substances also might enhance the ability of dogs to detect explosives, drugs, human remains, or other targets for which they are valued.

Keywords: Ecuador, hallucinogens, hunting, psychoactive plants, Shuar, Quichua

1. Introduction

Dogs (*Canis lupus familiaris*) in the New World originated from multiple Old World lineages that migrated with late Pleistocene humans across the Bering Strait (Leonard et al. 2002). They entered South America with the early human colonists and also were re-introduced by European explorers. Dogs were apparently absent in the Amazon Basin (until the historical period) but present in the Guyanas and the Orinoco River Basin. Following European contact, genetic evidence suggests that newly introduced European dog races began to replace native dogs throughout the Americas (Koster (2009). Hunting dogs are now common throughout much of the Amazon region (Fig. 1).



Fig. 1. Hunting dog in a Quichua village in Ecuador.

The role of dogs in human societies is diverse. They assist in warfare, detect odors, deter pest and predatory animals, guard property and people, guide the blind and deaf, protect other domesticated animals, provide human companionship, pull sleds, rescue lost and injured humans, and track and retrieve game animals. They also provide food and fur, serve as living blankets, and function in symbolic rituals (Diamond 1997, Coppinger and Schneider 1995, Hart 1995). Dogs play an important role in religions and rituals throughout the world. Ecuadorian Shuar believe that dogs are a gift from Nunkui, the earth mother (Bennett et al. 2002). According to the Quichua, dogs are gifts from sachahuarmi or sacharuna (forest spirits). They believe the canines protect hunters and family members from malevolent forest spirits called mal aire (bad air) and mal ojo (evil eye). They also believe that dogs dream and that they have souls (Kohn 2007). The Egyptian god Anbu (or Anubis) is often portrayed as a man with the

head of a dog or jackal (Gadalla 2001). Xolotl, twin of the Aztec god Quetzalcoatl, was the dog god and served as a guide to the dead (Fernández 1992). As part of a burial ritual, Aztec inhabitants of Anahuac killed a dog and laid it beside a human corpse. They believed that four years after death, the dog carried human soul to Chicunauhapan, the ultimate resting place of the dead (Beyer 1908). Dogs possess social-cognitive traits that allow them to communicate with humans in ways unlike any other animal (Hare et al. 2002).

In lowland areas of the Neotropics, the primary role of canines is to assist in hunting wild game. Hunting efficiency using dogs compares favorably to other forms of hunting (Koster 2009). The percentage of hunting trips that included dogs varies widely across cultures from a high of 83% (Mayangna and Miskito of Nicaragua) to 3% (Piro of Peru). Hunting success with dogs depends in large part on the targeted species. Although canines can be employed for any terrestrial species, they are particularly effective against pacas (*Cuniculus paca*, Fig. 2), agoutis (*Dasyprocta* spp.), and other animals that thrive in anthropogenic environments. The absence of dogs among some lowland cultures may be due to high mortality rates of dogs, rather than a canine aversion.



Figure 2. Spotted paca (*Cuniculus paca*) in Ecuador.

Mortality in Neotropical dogs results from the interaction of factors including hunting-induced wounds, malnutrition, microbial pathogens, and parasitic infections. Owing to their importance in hunting, it is not surprising that many cultures have a robust pharmacopoeia especially for dogs (e.g., Bennett et al. 2002; Lans et al. 2000, 2001; Leonard et al. 2002; Jernigan 2009). Nonetheless, ethnoveterinary medicinal research is incipient (Nobrega Alves et

al. 2010). Within many cultures, hunting dogs receive particularly good care (Koster 2009). A Shuar woman, for example, may nurse a pup along with her own children (Bennett et al. 2002). In training dogs, both the Shuar and Quichua maintain the animals with a minimal diet supplemented with wild plants. While many plant species are employed to target canine illnesses, the majority are used to enhance the hunting ability of dogs. In a study that focused exclusively on ethnoveterinary practices, Jernigan (2009) identified 34 plants, that the Peruvian Aguaruna give to their dogs, most often to improve their hunting prowess. Plants are employed in baths to reduce their scent or to mask odors and thus decreasing their detectability by the targeted prey. Plants also function to clean buccal and nasal cavities to enhance olfaction (e.g., Lans et al. 2001, Sanz-Biset et al. 2009) or to enhance night vision (Wilbert 1987).

Neotropical hunters employ magic, rituals, and charms to improve their hunting success and similar methods are used on dogs (Koster 2009, Shephard 2002). Koster (2009) notes the “occasional” use of hallucinogens, but the use psychoactive plants is actually frequent and widespread in many parts of the Old and, especially, the New World tropics (e.g., Bennett et al. 2002). The employment of psychoactive substances to enhance hunting ability seems to be counterintuitive, yet its prevalence suggests that it is both adaptive and that it has an underlying pharmacological explanation. In this paper, we address three questions:

1. What plants do the Ecuadorian Shuar and Quichua give to dogs to improve their hunting abilities?
2. What plants do other cultures give to dogs?
3. What is the likely pharmacological basis for the use of these plants, particularly the psychoactive ones?

The Shuar and Quichua are the largest indigenous groups in lowland Ecuador. They mostly reside at elevations from 300 to 1,200 m in terra firme forests. This territory spans two of Holdridge's (1967) life zones, tropical moist forest and premontane tropical wet forest. Study sites were located in the Provinces of Morona-Santiago and Napo (Fig. 3).

Both groups are horticulturists, growing manioc (*Manihot esculenta*) and plantains (*Musa × paradisiaca* L.) as their principle starches.

Hunting (Fig. 4) and fishing supplement animal sources of protein from domesticated chickens and pigs. More data on the research sites and the two cultures can be found in Bennett et al. (2002) and Bennett and Alarcón (1994).

2. Methods

The Shuar data analyzed here was published in Bennett (1992a, 1992b, 1994) and Bennett et al. (2002). The Quichua data comes from Alarcón (1988), Bennett and Alarcón (1994) and our unpublished field

notes. Voucher specimens are deposited in QCNE in Ecuador with duplicates in NY and MO in the U.S. We located data on ethnoveterinary medicine from other tropical cultures from ethnobotanical monographs, JEP publications, and searches using Web of Science and Google Scholar. Family circumscriptions and species names have changed since many of the data sources were first published. Data presented here follows APG III family circumscriptions (Angiosperm Phylogeny Group 2009). Species names follow The Plant List (2015) except as




Figure 3. Provinces of Morona-Santiago, where the Shuar, and Napo, where the Quichua, of this study reside.



Figure 4. Shuar hunter in Yukutais with blowgun.

noted. Author citations are included in the text for species not cited in Tables 1 or 2. Chemical data was found using PubMed and SciFinder.

3. Results

3.1 Shuar and Quichua

The Shuar and Quichua employ at least 22 species for dogs (Table 1). The studies from which these data were drawn did not focus on ethnoveterinary medicines. It is therefore likely that more exist. In most case, the plants have corresponding human uses. However, some species or varieties are especially designated for canines. Four Shuar ethnoveterinary plants carry the name yawá, which means dog in the Shuar language: yawá kunkunari (*Justicia pectoralis*), yawá urints (*Alternanthera paronychioides*), yawá piripiri (*Cyperus* sp.) and yawá maikua (*Brugmansia versicolor*). With the exception of *Brunfelsia grandiflora* D. Don., all the principal Shuar hallucinogens are given to dogs.

Seven of the plants were utilized for purely medical reasons, mostly to treat botfly or other infections. The remaining plants are given to hunting dogs specifically to enhance their hunting prowess. Nine were used for the general purpose of improving hunting ability. A mixture of manioc and akapmas (*Fittonia albivenis*) was said to improve the ability to track game. Kunápik (*Tabernaemontana sananho*) and yawá piripiri (*Cyperus* sp.) appear to initiate hunting predilections in dogs. Quichua give payanshi (*Abuta grandifolia*) to their hunting dogs to keep them quiet and both the Quichua and the Shuar give the potent stimulant wais (*Ilex guayusa*,



Figure 5. *Ilex guayusa* (Aquifoliaceae) leaves known as wais in the Shuar and guayusa in the Runa languages.

Fig. 5) to their hunting dogs so that “they will not be lazy.”

Accepted manuscript

Table 1. Shuar and Quichua plants given to dogs. Source of data: when not specified - Bennett et al. 2002; footnotes indicate other sources. Family names follow APG III (Angiosperm Phylogeny Group 2009). Species names follow The Plant List (2105), except where indicated. SH = Shuar, QU = Quichua, SP=Spanish. For plant parts: BK=bark, FL=flower, FR=fruit, IF=inflorescence, LF=leaf, LX=latex, TU=Tuber.

Species	Family	Voucher	Common Name	Use
<i>Fittonia albivenis</i> (Lindl. ex Veitch) Brummitt	Acanthaceae	Bennett 3712	AKAPMAS [SH: derived from akap "liver"]	SH give masticated LF, mixed with <i>Manihot esculenta</i> & meat, to improve dogs' ability to track game
<i>Justicia pectoralis</i> Jacq.	Acanthaceae	Pujupet 1009	YAWÁ KUNKUNARI [SH: yawá "dog" kunkunari unknown]	SH mix LF mixed with food, to improve dogs' hunting ability
<i>Alternanthera paronychioides</i> A. St.-Hil.	Amaranthaceae	Warush 44	YAWÁ URINTS [SH: yawá "dog" urints known"]	SH give LF to improve dogs' hunting ability
<i>Tabernaemontana sananho</i> Ruiz & Pav.	Apocynaceae	Bennett 4081	KUNÁPIK [SH: kunápiik <i>Tabernaemontana</i> sp.]	SH give LX to dogs so that "they won't be vagrants." BK also given to dogs that "don't hunt."
<i>Ilex guayusa</i> Loes.	Aquifoliaceae	Bennett 3659	WAIS [SH: waís from QU: guayusa = <i>Ilex guayusa</i>]	SH give LF decoction to hunting dogs
<i>Anthurium eminens</i> Schott	Araceae	Bennett 4543	TACOTA SHIPU [QU: toccata unknown shipu "infested with worms"]	QU grind the IN, then apply it to botfly infections in animals ¹
<i>Anthurium gracile</i> (Rudge) Schott ²	Araceae	Bennett 3705	WANKAT [SH: wankat <i>Anthurium</i> or <i>Philodendron</i> spp.]	QU apply FR to kill botfly larvae in cattle & dogs ²
<i>Caladium bicolor</i> (Aiton) Vent.	Araceae	Warush 50	USHU [SH: ushu = <i>Caladium bicolor</i>]	SH treat animals infested with worms with LF sap

<i>Caladium schomburgkii</i> Schott	Araceae	Utitiáj 16	APINIÚ WANCHÚP [SH: napiniú may be from napi "snake" wanchúp perhaps <i>Xanthosoma</i> sp.]	SH give plant extract to dogs to improve their hunting ability
<i>Cyperus</i> sp.	Cyperaceae	Warush 23	YAWÁ PIRIPIRI [SH: yawa "dog" piri-piri "Cyperaceae spp."]	SH give TU given to dogs or mix with saliva & place in dog's eyes to make them become hunters
<i>Euphorbia hirta</i> L.	Euphorbiaceae	Bennett 4010	LOMO PACHINA [perhaps SP: lomo "back" QU: pachina maybe from papachina= <i>Colocasia esculenta</i>]	QU give LX to dogs to make them better hunters ¹
<i>Manihot esculenta</i> Crantz	Euphorbiaceae	Bennett 3320	MAMA [SH: mama = <i>Manihot esculenta</i>]	SH give TU mixed with masticated <i>Fittonia albivenis</i> LF to dogs so that they can follow animal trails
<i>Desmodium</i> sp.	Fabaceae	Kunkumas 191	MÍIKMAN [SH: derived from miik "bean"]	SH bathe dogs with a LF decoction.
<i>Casearia</i> aff. <i>commersoniana</i> Cambess	Salicaceae	Bennett 4135	YAMAKAI [SH: also refers to <i>Ryania speciosa</i> Vahl var. <i>speciosa</i>]	SH treat mange with plant
<i>Salvia angulata</i> Benth.	Lamiaceae	Gómez 397	PURGA PERRO [SP: "dog purgative"]	SH use FL & FR decoction for a variety of ailments. Common names suggests its ethnoveterinary use
<i>Abuta grandifolia</i> (Mart.) Sandwith	Menispermaceae	Bennett 4400	PAYANSHI [QU: payanshi = <i>Abuta grandifolia</i>]	QU give BK decoction to dogs before hunting to keep them quiet ¹
<i>Ficus insipida</i> Willd. ssp. <i>insipida</i>	Moraceae	Anananch 154	JAPÁ WAMPÚ [SH: japá deer wampúch "light weight"]	SH give LX to dogs, probably as a anthelmintic

<i>Osteophloeum platyspermum</i> (Spruce ex A. DC.) Warb.	Myristicaceae	Palacios 4694	ANYA CASPI (QU: anya "counsel?" huapa "Myristicaceae spp.")	QU give resin to dogs to improve their hunting abilities. ³
<i>Virola duckei</i> A.C. Sm.	Myristicaceae	Palacios 1780	HUAPA BLANCA (QU: huapa "Myristicaceae spp." SP: blanca "white")	Employed in same manner as <i>Osteophloeum platyspermum</i> ³
<i>Brugmansia suaveolens</i> (Humb. & Bonpl. ex Willd.) Bercht. & C. Presl	Solanaceae	Bennett 3312	MAIKUA [SH: maikua = <i>Brugmansia</i> spp.] GUANDU [QU: maikua = <i>Brugmansia</i> spp.]	QU & SH give extracts to dogs to improve their hunting ability.
<i>Brugmansia versicolor</i> Lagerh.	Solanaceae	Shiki 333	YAWÁ MAIKUA [SH: yawá "dog" maikua <i>Brugmansia</i> spp.]	SH give 3 crushed LF every 12 hours to make them good hunters
<i>Brugmansia × insignis</i> (Barb. Rodr.) Lockwood ex E.W. Davis	Solanaceae	Bennett 4003	GUANDU LUMACHAG [QU: guandu = <i>Brugmansia</i> spp. lumachag "unknown"]	QU apply macerated LF to dogs' noses to make them better hunters ¹

¹Alarcón (1988), ²Bennett and Alarcón (unpublished field notes), ³Bennett and Alarcón (1994)

²Author citation follows TROPICOS 2015.

Table 2. Examples of other plants given to dogs to improve their hunting abilities. Family names follow APG III (Angiosperm Phylogeny Group 2009). Species names follow The Plant List (2105). Source of data: see footnotes. For plant parts: BK=bark, FL=flower, FR=fruit, IF=inflorescence, LF=leaf, LX=latex, PL=whole plant, RH=rhizome, SD=seed, ST = stem, TU=Tuber.

Species	Family	Common Name	Location	Use
<i>Aframomum melegueta</i> K. Schum.	Zingiberaceae	guinea pepper	Trinidad and Tobago ¹	Dried SD ground to a powder, then sprinkled on dog's food
<i>Anadenanthera colubrina</i> (Vell.) Brenan	Fabaceae	hujilca	Argentina ²	SD snuff given to dogs to make them more alert
<i>Anadenanthera peregrina</i> (L.) Speg.	Fabaceae	Yopo	Colombia ²	SD snuff given to dogs to make them more alert
<i>Ardisia</i> cf. <i>denhamioides</i> S. Moore	Primulaceae	sunun	West Papua ³	LF fed to hunting dogs to improve their performance.
<i>Aristolochia rugosa</i> Lam.	Aristolochiaceae	mat	Trinidad and Tobago ¹	PL decoction used to bathe lazy dogs
<i>Banisteriopsis caapi</i> (Spruce ex Griseb.) C. V. Morton	Malpighiaceae	-	Peru ⁴	See <i>Couroupita guianensis</i> [A]
<i>Brugmansia suaveolens</i> (Humb. & Bonpl. ex Willd.) Bercht. & J. Presl	Solanaceae	-	Peru ⁴	See <i>Couroupita guianensis</i> [A]
<i>Brugmansia</i> sp.	Solanaceae	lumu cuchi huandu	Ecuador (Quichua) ⁵	See <i>Tabernaemontana sananho</i> [A]
<i>Caladium bicolor</i> (Ait.) Vent.	Araceae	-	Ecuador (Kofan) ¹³	LF placed in dogs nostril to make them better hunters for wild pigs.
<i>Caladium</i> sp.	Araceae	ushu	Peru (Aguaruna) ⁶	LF fed to dogs
<i>Calliandra angustifolia</i> Spruce ex Benth.	Fabaceae	-		See <i>Couroupita guianensis</i> [A]
<i>Capsicum annuum</i> L.	Solanaceae	bird pepper	Trinidad and Tobago ¹	Juice from 2 small FR placed in dog's nose

<i>Casearia negrensis</i> Eichler	Salicaceae	ituchi runtu	Peru ^{4,7}	so it find game & follow scent BK & LF given to dogs to prepare them for hunting
<i>Cecropia peltata</i> L.	Urticaceae	bois canôt	Trinidad and Tobago ¹	Dry LF is put in water with <i>Jatropha gossypifolia</i> . Water left open for nine days until larvae are seen, then used to bathe dog.
<i>Colocasia esculenta</i> (L.) Schott [possibly]	Araceae	Bolobolo	Papua New Guinea ⁸	Unspecified part fed to dogs to make them wild, aggressive & sensitive for hunting wild pig
<i>Couroupita guianensis</i> Aubl. [A]	Lecythidaceae	aya uma	Peru ⁴	Depurative decoction including <i>Banisteriopsis caapi</i> , <i>Psychotria viridis</i> , <i>P. carthagenensis</i> , <i>Brugmansia suaveolens</i> , <i>Calliandra angustifolia</i> , <i>Tovomita</i> aff. <i>stylosa</i> & <i>Zygia longifolia</i> .
<i>Couroupita guianensis</i> Aubl. [B]	Lecythidaceae	ayahúma	Peru ⁹	Unspecified part given to dogs to make them stronger & to increase their hunting abilities
<i>Couroupita subsessilis</i> Pilg.	Lecythidaceae	shishim	Peru (Aguaruna) ⁶	BK & LF inhaled in mouth or nose to improve hunting ability
<i>Croton gossypifolius</i> Vahl	Euphorbiaceae	blood bush	Trinidad and Tobago ¹	Dog bathed in a LF decoction along <i>Petiveria alliacea</i> RT & <i>Renalmia alpina</i> LF & RT if dog is not performing as well as in past
<i>Cyperus</i> sp.	Cyperaceae	yawaa pijipij	Peru (Aguaruna) ⁶	RT fed to dogs to improve their hunting ability
<i>Cyrtocymura scorpioides</i> (Lam.) H. Rob.	Asteraceae	ruckshun	Trinidad and Tobago ¹	Dogs bathed with a LF decoction so that

<i>Dendrobium pulchellum</i> Roxb. ex Lindl.	Orchidaceae	-	Indochina ¹⁰	they will be more alert FL fed to dogs to make them better hunters
<i>Dendrobium</i> spp	Orchidaceae	-	Solomon Islands ¹¹	FL, resembling dog heads, fed to hunting dogs to increase their courage in the chase
<i>Dendropanax arboreus</i> (L.) Decne. & Planch.	Araliaceae	fei jein	Trinidad and Tobago ^{1,7}	Combined with LF of <i>Monstera dubia</i> , <i>Siparuna guianensis</i> <i>Solanum</i> spp. & <i>Syngonium podophyllum</i> to bathe dogs
<i>Dracontium</i> sp	Araceae	uchi santanik	Peru (Aguaruna) ⁶	RT fed to dog to improve hunting ability
<i>Eschweilera subglandulosa</i> (Steud. ex O. Berg) Miers	Lecythidaceae	guatacare	Trinidad and Tobago ^{1,7}	see <i>Piper marginatum</i>
<i>Jatropha curcas</i> L.	Euphorbiaceae	white physic nut	Trinidad and Tobago ¹	3 LV mixed with 3 LF of <i>Jatropha gossypifolia</i> , crushed then put in water to bathe dog
<i>Jatropha gossypifolia</i> L.	Euphorbiaceae	red physic nut	Trinidad and Tobago ¹	See <i>Jatropha curcas</i>
<i>Mansoa alliacea</i> (Lam.) A.H. Gentry	Bignoniaceae	ajo sacha	Peru ⁴	Stem & root BK macerated with with <i>Petiveria alliacea</i> as a body odor modifying agent for hunting & fishing
<i>Mansoa</i> sp.	Bignoniaceae	kaep	Peru (Aguaruna) ⁶	LF, RT, BK & ST tips inhaled in mouth, nose, or fed to dogs to improve their ability to hunt

<i>Momordica charantia</i> L.	Cucurbitaceae	caraaili	Trinidad and Tobago ^{1,7}	PL used to bathe dogs so that it will catch agoutis. See <i>Dendropanax arboreus</i>
<i>Monstera dubia</i> (Kunth) Engl. & K. Krause	Araceae	sei jein	Trinidad and Tobago ^{1,7}	RT inhaled in mouth & nose to improve hunting ability
<i>Nectandra cuneatocordata</i> Mez	Lauraceae	mantaga	Peru (Aguaruna) ⁶	LF used to cleans dog's nose to improve its ability to follow a scent
<i>Nicotiana tabacum</i> L. [A]	Solanaceae	tobacco	Trinidad and Tobago ¹	Mixture of tobacco & <i>Zingiber officinale</i> applied to eyes to improve night vision. See <i>Tabernaemontana sananho</i> [A]
<i>Nicotiana tabacum</i> L. [B]	Solanaceae	tobacco	South America ¹¹	Dogs bathed with ground RT to make them more alert
<i>Nicotiana tabacum</i> L. [C]	Solanaceae	-	Ecuador (Runa) ⁵	Pieces of LF fed to hunting dogs to improve their performance
<i>Petiveria alliacea</i> L.	Phytolaccaceae	kojo root	Trinidad and Tobago ¹	FR given to bathe dogs for "cross"
<i>Phrynium</i> sp.	Marantaceae	asin	West Papua ³	LF used to bathe dogs
<i>Phyllanthus urinaria</i> L.	Phyllanthaceae	-	Trinidad and Tobago ¹	LF used to bathe dogs so the will catch agoutis. Some hunters add <i>Eschweilera subgladulosa</i> LV.
<i>Piper hispidum</i> Sw.	Piperaceae	candle bush	Trinidad and Tobago ¹	Crushed ST & LV or RT put in dog's nose or dog's nose washed with a solution made from those parts of the plant
<i>Piper marginatum</i> Jacq.	Piperaceae	agouti bush	Trinidad and Tobago ^{1,7}	Crushed LF used to bathe dog for "cross"
<i>Piper ovatum</i> Vahl	Piperaceae	pot bush	Trinidad and Tobago ¹	RT & LF inhaled in mouth or nose or fed to dogs to improve their hunting ability
<i>Piper peltatum</i> L.	Piperaceae	sun bush	Trinidad and Tobago ¹	
<i>Piper</i> sp. 1	Piperaceae	ampagpag	Peru (Aguaruna) ⁶	

<i>Piper</i> sp. 2	Piperaceae	shishig	Peru (Aguaruna) ⁶	LF fed to dogs to improve their hunting ability
<i>Pithecellobium unguis-cati</i> (L.) Benth.	Fabaceae	cat's claw	Trinidad and Tobago ^{1,7}	Given to dogs so that they will follow game tenaciously
<i>Psychotria carthagenensis</i> Jacq	Rubiaceae	-	Peru ⁴	See <i>Tabernaemontana sananho</i> [A]
<i>Psychotria viridis</i> Ruiz & Pav.	Rubiaceae	-	Peru ⁴	See <i>Tabernaemontana sananho</i> [A]
<i>Renalmia alpina</i> (Rottb.) Maas	Zingiberaceae	mardi gras	Trinidad and Tobago ^{1,7}	5-7 shoots of are pounded & put in water to bathe dog; FR attract lice, dog will closely pursue
<i>Saccharum officinarum</i> L.	Poaceae	sugar cane	Trinidad and Tobago ^{1,7}	FR & LF used to bathe dogs. Deer eat LF, therefore dog can track deer
<i>Sarcoglottis metallica</i> (Rolfe) Schltr.	Orchidaceae	lappe bush	Trinidad and Tobago ^{1,7}	PL used to bathe dog so that it will track lappe
<i>Schuurmansia henningsii</i> K. Schum.	Ochnaceae	sembererk	West Papua ³	SD & LF fed to dogs to improve their performance
<i>Siparuna guianensis</i> Aubl.	Monimiaceae	dead man's bush	Trinidad and Tobago ^{1,7}	Ssee <i>Dendropanax arboreus</i>
<i>Solanum</i> spp.	Solanaceae	Devil pepper	Trinidad and Tobago ^{1,7}	Ssee <i>Dendropanax arboreus</i>
<i>Syngonium podophyllum</i> Schott	Araceae	Matapal-kit	Trinidad and Tobago ^{1,7}	See <i>Dendropanax arboreus</i>
<i>Tabernaemontana sananho</i> Ruiz & Pav. [A]	Apocynaceae	tsita	Ecuador (Runa) ⁵	Mixture given to hunting dogs so that they can communicate with their masters
<i>Tabernaemontana sananho</i> Ruiz & Pav. [B]	Apocynaceae	kunakip	Peru (Aguaruna) ⁶	BK & RT inhaled in mouth or nose or fed to dogs to improve hunting ability
<i>Tabernaemontana sananho</i> Ruiz & Pav. [C]	Apocynaceae	-	Ecuador (Secoya) ¹³	LX from FR applied to dogs nose so that they can "smell far" in huntingatex

<i>Tovomita</i> aff. <i>stylosa</i> Hemsl.	Clusiaceae	-	Peru ⁴	See <i>Couroupita guianensis</i> [A]
<i>Xanthosoma brasiliense</i> (Desf.) Engl.	Araceae	hog tannia	Trinidad and Tobago ^{1,7}	Hogs eat tubers. Ground tuber sprinkled on dog's food so it can track hogs. See <i>Xanthosoma brasiliense</i>
<i>Xanthosoma undipes</i> (K. Koch & C.D. Bouché) K. Koch	Araceae	hog tannia	Trinidad and Tobago ^{1,7}	LF given to dogs so that they will walk straight
<i>Xiphidium caeruleum</i> Aubl.	Haemodoraceae	walk fast	Trinidad and Tobago ^{1,7}	
<i>Zingiber officinale</i> Roscoe [A]	Zingiberaceae	ginger	South America ¹²	See <i>Nicotiana tabacum</i>
<i>Zingiber officinale</i> Roscoe [B]	Zingiberaceae	hinde	West Papua ¹⁰	RH & LF are fed to hunting dogs to improve their performance
<i>Zygia longifolia</i> (Willd.) Britton & Rose	Fabaceae	-	Peru ⁴	See <i>Couroupita guianensis</i> [A]
not determined	Lamiaceae	chiujip	Peru (Aguaruna) ⁶	LF fed to dogs to improve their hunting ability
not determined	Zingiberaceae	chiag	Peru (Aguaruna) ⁶	LF inhaled in mouth to improve hunting ability
not determined	-	turpentine bush	Trinidad and Tobago ¹	Dogs bathed with crushed LF
not determined	-	Goma'uwe	Papua New Guinea ⁸	Unspecified part fed to hunting dogs to improve their sense of smell
not determined	Acanthaceae	tumbaik	Peru (Aguaruna) ⁶	RT & LF fed to dogs to improve hunting ability
not collected	-	chijum	Peru (Aguaruna) ⁶	LF fed dogs to improve hunting ability

¹Lans et al. (2001), ²Schultes (1998), ³Miliken (1999), ⁴Sanz-Biset et al. (2009), ⁵Kohn (2007), ⁶Jernigan (2009), ⁷Use attributed to Doctrine of Signatures, but see Bennett (2007 & 2008) for a discussion of the doctrine, ⁸Flavelle (1991), ⁹McKenna et al. (1995), ¹⁰Bennett (1995), ¹¹Anonymous (2009a), ¹²Wilbert (1987), ¹³Schultes and Raffauf (1990)

3.2 Other Cultures

The use of plants to improve the hunting ability is best documented in Ecuador and Peru but examples can be found in other South American countries as well as the Caribbean, Indochina, Papua New Guinea, and the Solomon Islands (Table 2). Examples from the literature revealed 71 citations of 65 species that are used in 54 combinations. Of these, the majority (43) are said to improve hunting ability (e.g., *Dendrobium pulchellum*). Five enhance hunting success for specific game (e.g., *Xanthosoma brasiliense* for wild hogs). Four are believed to make hunting dogs more alert (e.g., *Petiveria alliacea*) and four are said to specifically enhance olfaction. The Secoya of Ecuador apply latex from *Tabernaemontana sananho* fruits to a dog's nose so that "it can smell far." A mixture of ginger (*Zingiber officinale*) and tobacco (*Nicotiana tabacum*, Fig. 6) is thought to enhance night vision in both hunters and their dogs. Ten plants are employed in baths for hunting dogs. A mixture that includes *Tabernaemontana sananho* and *Brugmansia* sp. is given to dogs so that they "can communicate with their masters."



Figure 6. *Nicotiana tabacum* (Solanaceae) plug used by a Quichua shaman in healing ceremonies.

3.3 Plants to Improve Hunting Ability

The combined data from the Shuar and Quichua data (Table 1) and the literature (Table 2) omitting those species that are not related directly to hunting or those species that have not been determined to at least the genus, reveals 71 species in 34 families that are given to dogs to improve their hunting ability (Table 3). There is some chemical data for most of the species or from close relatives. By combining the phytochemical data with the ethnobotanical reports of plants use, we classified each species into a likely pharmacological category. Twenty six are

Table 3. Combined data from Tables 1 and 2 minus those species that have an explicitly ethnoveterinary use and are not related directly to hunting or those species that have not been determined to at least the genus. Family plus number of citations for family, species plus number of citations for genus/species). Activity column lists the probable activity in dog based on the plants use and chemistry. Chemistry column lists the compounds likely responsible for the activity.

Family	Species	Activity	Chemistry
Acanthaceae (2)	<i>Fittonia albivenis</i> (1/1) <i>Justicia pectoralis</i> (1/1)	psychoactive - hallucinogenic (?) psychoactive - hallucinogenic	dimethyltryptamine (?) betaine, coumarin & umbelliferone ¹ ; reported to have dimethyltryptamine (DMT)
Amaranthaceae (1)	<i>Alternanthera paronychioides</i> (1/1)	unknown	unknown - inhibits xanthine oxidase ²
Apocynaceae (4)	<i>Tabernaemontana sananho</i> (1/4)	psychoactive - hallucinogenic (?)	unknown, indole alkaloids from <i>Tabernaemontana holstii</i> ³
Aquifoliaceae (1)	<i>Ilex guayusa</i> (1/1)	psychoactive - stimulant	caffeine & other methylxanthine alkaloids ⁴
Araceae (9)	<i>Caladium bicolor</i> (3/1) <i>Caladium schomburgkii</i> (3/1) <i>Caladium</i> sp. (3/1) <i>Colocasia esculenta</i> (1/1) <i>Dracontium</i> sp. (1/1) <i>Monstera dubia</i> (1/1) <i>Syngonium podophyllum</i> (1/1) <i>Xanthosoma brasiliense</i> (2/1) <i>Xanthosoma undipes</i> (2/1)	olfactory sensitizer olfactory sensitizer olfactory sensitizer olfactory sensitizer olfactory sensitizer olfactory sensitizer olfactory sensitizer olfactory sensitizer	calcium oxalate crystals ⁵ calcium oxalate crystals ⁵ calcium oxalate crystals ⁵ calcium oxalate crystals ^{5,6,7} calcium oxalate crystals ^{5,6,7} calcium oxalate crystals ^{5,6,7} calcium oxalate crystals ^{5,6,7} calcium oxalate crystals ^{5,6,7}
Araliaceae (1)	<i>Dendropanax arboreus</i> (1/1)	depuarative, deodorant, irritant	falcarinol, dehydrofalcarinol, undetermined diynene, falcarindiol, dehydrofalcarindiol, polyacetylenes (dendroarboreols A and B) ^{8,9}
Aristolochiaceae (1)	<i>Aristolochia rugosa</i> (1/1)	depuarative, anti-inflammatory, antimicrobial	aristolochic acid ¹⁰ (?)
Asteraceae (1)	<i>Cyrtocymura scorpioides</i> (1/1)	anti-tumor, vulnerary	sesquiterpene lactones ¹¹

Bignoniaceae (2)	<i>Mansoa alliacea</i> (2/1)	depuarive, deodorant	allyl polysulfides ¹²	
	<i>Mansoa</i> sp. (2/1)	depuarive, deodorant	likely similar to <i>Mansoa alliacea</i>	
Clusiaceae (1)	<i>Tovomita</i> aff. <i>stylosa</i> (1/1)	depuarive, deodorant, antimicrobial	xanthones likely ¹³	
Cucurbitaceae (1)	<i>Momordica charantia</i> (1/1)	depuarive, deodorant, antimicrobial	momordicin, cucurbitacin b & other triterpenoid saponins ¹⁴	
Cyperaceae (2)	<i>Cyperus</i> sp. (2/1)	psychoactive – hallucinogenic	ergot alkaloids ¹⁵	
	<i>Cyperus</i> sp. (2/1)	psychoactive – hallucinogenic	ergot alkaloids ¹⁵	
Euphorbiaceae (5)	<i>Croton gossypifolius</i> (1/1)	depuarive, deodorant, antimicrobial	essential oil dominated by oxygenated sesquiterpenes including alpha-cedrene oxide, spathulenol, valencene, geranyl-pentanoate, alpha-cadinol, germacrene d & longifolene) ¹⁷	
	<i>Euphorbia hirta</i> (1/1)	antimicrobial	flavonols including euphorbins, kaempferol, myricetin, quercetin & rutin ¹⁶	
	<i>Jatropha curcas</i> (2/1)	depuarive, deodorant, antimicrobial	hydrogen cyanide, jatrophine ^{18,19}	
	<i>Jatropha gossypifolia</i> (2/1)	depuarive, deodorant, antimicrobial	cyclic octapeptide (cyclogossine b) & cyclic heptapeptide (cyclogossine a) ²⁰ , diterpene jatrophene ²¹	
Fabaceae (5)	<i>Manihot esculenta</i> (1/1)	carrier	not applicable	
	<i>Anadenanthera colubrina</i> (2/1)	psychoactive – hallucinogenic	buforenine, n-n-dimethyltryptamine ¹³	
	<i>Anadenanthera peregrina</i> (2/1)	psychoactive - hallucinogenic	buforenine, n-n-dimethyltryptamine ¹³	
	<i>Calliandra angustifolia</i> (1/1)	psychoactive – hallucinogenic (?), stimulant	amino acids ²² , reported to contain tetrahydroharmaline	
	<i>Pithecellobium unguis-cati</i> (1/1)	psychoactive – hallucinogenic (?), stimulant	unknown	
	<i>Zygia longifolia</i> (1/1)	depuarive, deodorant, antimicrobial (?)	volatile isoprenes, but poorly known ²³	
Haemodoraceae (1)	<i>Xiphidium caeruleum</i> (1/1)	antimicrobial (?)	phenylphenalenone-type compounds ²⁴	

Lauraceae (1)	<i>Nectandra cuneatocordata</i> (1/1)	unknown	unknown, genus contains alkaloids & terpenes ¹³ & lignans, neolignans, & lignoids ^{25,26,27}
Lecythidaceae (4)	<i>Couroupita guianensis</i> (2/3)	depuartive, deodorant, antimicrobial, antiinflammatory	tryptanthrin ²⁸
	<i>Eschweilera subglandulosa</i> (1/1)	depuartive, deodorant, antimicrobial	ellagic acid derivatives ²⁹ ; triterpenes ³⁰
Malpighiaceae (1)	<i>Banisteriopsis caapi</i> (1/1)	psychoactive – hallucinogenic	beta-carboline alkaloids (harmine, harmaline & tetrahydroharmine) ^{13,61,62}
Marantaceae (1)	<i>Phrynium</i> sp. (1/1)	unknown	unknown, rosmarinic acid, chlorogenic acid & rutin reported from family ³¹
Menispermaceae (1)	<i>Abuta grandifolia</i> (1/1)	psychoactive – sedative (?)	tropoloisoquinoline ³² & bisbenzylisoquinoline ³³ alkaloids
Monimiaceae (1)	<i>Siparuna guianensis</i> (1/1)	depuartive, deodorant	oxoaporphine alkaloid liriodenine, oxidized derivative of β -elemene-curzerenone & phenylpropanoids (myristicin & eugenol methyl ether) ³⁴
Myristicaceae (2)	<i>Osteophloeum platyspermum</i> (1/1)	psychoactive – hallucinogenic	unknown, presumably DMT
	<i>Virola duckei</i> (1/1)	psychoactive – hallucinogenic	unknown, presumably DMT
Myrsinaceae (1)	<i>Ardisia</i> cf. <i>denhamioides</i> (1/1)	Unknown	unknown - triterpenoid saponins common in the genus ³⁵
Ochnaceae (1)	<i>Schuurmansia henningsii</i> (1/1)	psychoactive – stimulant (?)	alkaloids ³⁶
Orchidaceae (3)	<i>Dendrobium pulchellum</i> (2/1)	psychoactive – hallucinogenic (?)	unknown, dendrobine & other alkaloids reported from <i>D. nobile</i> ³⁷
	<i>Dendrobium</i> spp. (1/1)	psychoactive – hallucinogenic (?)	unknown, dendrobine & other alkaloids reported from <i>D. nobile</i> ³⁷
	<i>Sarcoglottis metallica</i> (1/1)	depuartive, deodorant	unknown, flavonoids ³⁸ & prenylated coumarins ³⁹ found in the closely related genus <i>Spiranthes</i>
Phyllanthaceae (1)	<i>Phyllanthus urinaria</i> (1/1)	depuartive	alcohol (triacotanol), phenolic acid (gallic acid), coumarins (including ellagic acid), flavonoids (astragalin, quercetin, quercitrin, isoquercitrin, rutin, kaempferol), sterols (including β -sitosterol), triterpenes (lupeol acetate & β -amyrin) ⁴⁰
Phytolaccaceae (1)	<i>Petiveria alliacea</i> (1/1)	depuartive, deodorant	benzaldehyde, benzoic acid, coumarin, trithiolaniacine ⁴¹ , cysteine

Piperaceae (6)	<i>Piper hispidum</i> (6/1)	depuartive, deodorant	sulfoxide derivatives ⁴² amide ⁴³ ; butenolides ⁴⁴ ; prenylated benzoic acid derivatives ⁴⁵ ; oxygenated sesquiterpenes & sesquiterpenes hydrocarbons including trans-nerolidol, caryophyllene oxide, beta-elemene, trans- beta-caryophyllene, curzerene, & germacrene b ⁴⁶ 39 components in essential oil including γ-terpinene, δ-elemene, α- copaene, β-elemene, β-caryophyllene, α-humulene, γ-elemene, 3,4- methylenedioxypropiphenone, and elemicin ⁴⁷ but highly variable ⁴⁸ ; propiophenones ⁴⁹ ; flavonoids including marginatoside ⁵⁰ delta-amorphene, cis-murola-4(14),5-diene, & gamma- murolene51, amides (piperovatine & piperlonguminine) ⁵² 4-nerolidylcatechol ^{53,53} ; aristolactams alkaloids (piperumbellactams a-d & n-hydroxyaristolam & n-p-coumaroyl tyramine ⁵⁵ unknown unknown
	<i>Piper marginatum</i> (6/1)	depuartive, deodorant, antimicrobial	
	<i>Piper ovatum</i> (6/1)	depuartive, deodorant, antimicrobial, anti-inflammatory	
	<i>Piper peltatum</i> (6/1)	depuartive, deodorant, antimicrobial, anti-inflammatory	
	<i>Piper</i> sp. (6/1)	depuartive, deodorant, olfactory sensitizer	
	<i>Piper</i> sp. (6/1)	depuartive, deodorant, olfactory sensitizer	
Poaceae (1)	<i>Saccharum officinarum</i> (1/1)	depuartive, deodorant, antiinflammatory	fatty acids ⁵⁶ ; flavone glycosides vitexin, orientin, luteolin-8-c- (rhamnosylglucoside), 4',5'-dimethyl-luteolin-8-c-glycoside, isomeric pair schaftoside-isoschaftoside, o-glycosides tricrin-7-o- neohesperidoside & tricrin-7-o-glycoside ⁵⁷ ; flavones (tricrin-7-o-beta- (6''-methoxycinnamic)-glucoside & orientin ⁵⁸ ; flavones (apigenin, luteolin & tricrin derivatives), phenolic acids (hydroxycinnamic, caffeic & sinapic acids) ⁵⁹ DMT commonly reported ¹⁴ but not found in one study ⁶⁰ ; triterpene (beta-sitosterol & ursolic acid) ⁶¹ DMT ^{62,63} but variable
Rubiaceae (2)	<i>Psychotria carthagenensis</i> (2/1)	psychoactive – hallucinogenic	
	<i>Psychotria viridis</i> (2/1)	psychoactive – hallucinogenic	

Salicaceae (1)	<i>Casaria negrensis</i> (1/1)	unknown	unknown, clerodane diterpenoids ⁶⁴ reported for genus
Solanaceae (11)	<i>Brugmansia</i> × <i>insignis</i> (6/1)	psychoactive – hallucinogenic	tropane alkaloids (atropine & scopolamine) ¹⁴
	<i>Brugmansia</i> sp. (6/1)	psychoactive – hallucinogenic	tropane alkaloids (atropine & scopolamine) ¹⁴
	<i>Brugmansia suaveolens</i> (6/2)	psychoactive – hallucinogenic	tropane alkaloids (atropine & scopolamine) ¹⁴
	<i>Brugmansia versicolor</i> (6/1)	psychoactive – hallucinogenic	tropane alkaloids (atropine & scopolamine) ¹⁴
	<i>Brugmansia</i> sp. (6/1)	psychoactive – hallucinogenic	tropane alkaloids (atropine & scopolamine) ¹⁴
Urticaceae (1)	<i>Capsicum annuum</i> L. (1/1)	olfactory sensitizer	capsiacins ^{14,65}
	<i>Nicotiana tabacum</i> (3/3)	psychoactive – stimulant, MAO inhibitor, ophthalmic	nicotine ^{14,66}
	<i>Solanum</i> spp. (1/1)	depuartive	Unknown
Zingiberaceae (4)	<i>Cecropia peltata</i> (1/1)	depuartive	chlorogenic acid & isoorientin ⁶⁷
	<i>Aframomum melegueta</i> (1/1)	depuartive, antimicrobial	CYP inhibitor ⁶⁸ ; 27 compounds sesquiterpene hydrocarbons (humulene & caryophyllene & their oxides) 17 monoterpenes ⁶⁹
	<i>Renealmia alpinia</i> (1/1)	depuartive, antimicrobial	labdane diterpenoids ⁷⁰ , triacylglycerols (including oleic, palmitic & palmitoleic acids), sterols, methylsterols & triterpenic alcohols ⁷¹ ;
			monoterpenes (b-pinene, limonene & b-phellandrene), b-carotene, nerolidol & manool, labdadiene-15,16-dial (i) ⁷¹
	<i>Zingiber officinale</i> (2/2)	ophthalmic	phenols (gingerols & shogaols), volatile oils (sesquiterpenes - β-bisabolene, (-)-zingiberene, β-sesquiphellandrene, & (+)-ar-curcumene; monoterpenes - geranial & neral) ⁷³

¹Macrae & Towers (1984), ²Chen et al. (2009), ³Kingston et al. (2006), ⁴Lewis et al. (1991), ⁵Oscarsson & Savage (2007), ⁶Savage et al. (2008), ⁷Prychid & Rudall (1999), ⁸Bernart et al. (1999), ⁹Hansen & Boll (1986), ¹⁰Heinrich et al. (2009), ¹¹Buskuhl et al. (2010), ¹²Zoghbi et al. (2009), ¹³Schultes & Raffaui (1990), ¹⁴Fatope et al. (1990), ¹⁵Plowman et al. (1990), ¹⁶Kumar et al. (2010), ¹⁷Suárez et al. (2011), ¹⁸Duke (1993), ¹⁹Thomas et al. (2008), ²⁰Auvin-Guette et al. (1997), ²¹Ravindranath, et al. (2003), ²²McKenna et al. (1995), ²³Geron et al. (2002), ²⁴Opitz & Schneider (2002), ²⁵Carvalho et al. (1986), ²⁶Moro et al. (1986), ²⁷Barbosa-Filho et al. (1989), ²⁸Pinheiro et al. (2010), ²⁹Yang et al. (1998), ³⁰Costa & Carvalho (2003), ³¹Abdullah et al. (2008), ³²Menachery & Cava (1980), ³³Steele et al. (1999), ³⁴Leitão et al. (2011), ³⁵Liu et al. (2011), ³⁶Pelletier (1996), ³⁷Southon & Buckingham (1989), ³⁸Donget al. (2008), ³⁹Peng et al. (2008), ⁴⁰Calixto et al. (1998), ⁴¹Anonymous (2009b), ⁴²Kubec & Musah (2001), ⁴³Alécio et al (1998), ⁴⁴Michel et al. (2010), ⁴⁵Friedrich et al. (2005), ⁴⁶Pino et al. (2009), ⁴⁷Ramos et al. (1986), ⁴⁸Andrade et al. (2008), ⁴⁹de Diaz & Gottlieb (1979), ⁵⁰Tillequin et al. (1978), ⁵¹Silva et al. (2009), ⁵²Rodrigues et al. (2008),

- ⁵³Kijjoo et al. (1980), ⁵⁴Núñez et al. (2005), ⁵⁵Tabopda et al. (2008), ⁵⁶Ledón et al. (2003), ⁵⁷Colombo et al. (2006), ⁵⁸Duarte-Almeida et al. (2007), ⁵⁹Duarte-Almeida et al. (2006),
⁶⁰Leal & Elisabetsky (1996), ⁶¹Lopes et al. (2000), ⁶²Pires et al. (2009), ⁶³Callaway et al. (2005), ⁶⁴Chen et al. (2008), ⁶⁵Cichewicz & Thorpe (1996), ⁶⁶Wilbert (1987), ⁶⁷Andrade-
Cetto & Cárdenas Váz (2010), ⁶⁸Agbonon et al. (2010), ⁶⁹Ajaiyeoba & Ekundayo (1999), ⁷⁰Zhou et al. (1997), ⁷¹Lognay et al. (1989), ⁷²Lognay et al. (1991), ⁷³Blumenthal et al.
(2000)

Accepted manuscript

depuratives/deodorants (e.g., *Siparuna guianensis*), and many of these also have antimicrobial or anti-inflammatory activity. Ten species are classified as olfactory sensitizers Araceae (e.g., *Caladium schomburgkii*). The largest category was psychoactives, with 25 species. Nineteen of these species are hallucinogens (e.g., *Banisteriopsis caapi*, Fig. 7) and most of the remaining are stimulants (e.g., *Ilex guayusa*). Two are ophthalmic (discussed previously). The remaining are either unknown or difficult to classify.



Figure 7. *Banisteriopsis caapi* (Malpighiaceae), known as natem in the Shuar and ayahuasca in the Runa languages, split stems.

3.3.1 Depuratives/Deodorants

More than half of the depuratives/deodorants have noticeably strong odors (Table 4). *Dendropanax arboreus* has a distinctive odor due to the presence of polyactetylenes. The specific epithets of *Mansoa alliacea* and *Petiveria alliacea*, together with some of their common names, refer to the plants' garlic-like odor. *Siparuna* and *Piper* spp. possess abundant volatile terpenoids compounds that contribute to their strong and distinctive aromas. Cucurbitacins found in *Momordica charantia* produce its characteristic and pungent smell. Sesquiterpenes and monoterpenes in *Renealmia alpinia* and *Zingiber officinale* contribute to the distinctive ginger aroma of these plants.

Table 4. Species, arranged by family, that are used as depuratives or deodorants.

Family	Species
Araliaceae	<i>Dendropanax arboreus</i>
Aristolochiaceae	<i>Aristolochia rugosa</i>

Bignoniaceae	<i>Mansoa alliacea</i>
	<i>Mansoa</i> sp.
Clusiaceae	<i>Tovomita</i> aff. <i>stylosa</i>
Cucurbitaceae	<i>Momordica charantia</i>
	<i>Croton gossypiifolius</i>
	<i>Jatropha gossypiifolia</i>
Fabaceae	<i>Zygia longifolia</i>
Lecythidaceae	<i>Couroupita guianensis</i>
	<i>Eschweilera subglandulosa</i>
Monimiaceae	<i>Siparuna guianensis</i>
Orchidaceae	<i>Sarcoglottis metallica</i>
Phyllanthaceae	<i>Phyllanthus urinaria</i>
Phytolaccaceae	<i>Petiveria alliacea</i>
Piperaceae	<i>Piper hispidum</i>
	<i>Piper marginatum</i>
	<i>Piper ovatum</i>
	<i>Piper peltatum</i>
	<i>Piper</i> sp.
	<i>Piper</i> sp.
Poaceae	<i>Saccharum officinarum</i>
	<i>Solanum</i> spp.
Urticaceae	<i>Cecropia peltata</i>
Zingiberaceae	<i>Aframomum melegueta</i>
	<i>Renealmia alpinia</i>

3.3.2 Olfactory Sensitizers

The olfactory sensitizers are dominated by the Araceae (Table 5). This family is characterized by the presence of irritating calcium oxalate crystals. Two other plants were

classified in this category *Capsicum annuum* with irritating capsaicins and a *Piper* sp. with unknown chemical components. In addition to applying *Capsicum annuum* around the eyes of their hunting dogs to enhance vision, the Quichua believe that this practice protects the animals from evil spirits.

Table 5. Species, arranged by family, that are used as olfactory sensitizers.

Family	Species
Araceae	<i>Caladium bicolor</i>
	<i>Caladium schomburgkii</i>
	<i>Caladium</i> sp.
	<i>Colocasia esculenta</i>
	<i>Dracontium</i> sp.
	<i>Monstera dubia</i>
	<i>Syngonium podophyllum</i>
	<i>Xanthosoma brasiliense</i>
	<i>Xanthosoma undipes</i>
Piperaceae	<i>Piper</i> sp.
Solanaceae	<i>Capsicum annuum</i> L.

3.3.3 Psychoactives

The psychoactive plants given to dogs are dominated by hallucinogens (Table 6). While most of these are well-known as hallucinogenic plants and are commonly used in shamanistic rituals (e.g., *Anadenanthera peregrina*, *Banisteriopsis caapi*) the activity of others is yet to be determined (e.g., *Fittonia albivenis*, *Dendrobium pulchellum*). The stimulants *Ilex guayusa* (caffeine & other methylxanthine alkaloids) and *Nicotiana tabacum* also are given to hunting

dogs. Quichua heat tobacco leaves, then administer them through the noses of their dogs to keep them active and resilient during the hunting trips.

Table 6. Species, arranged by family, that have probable psychoactive effects.

Family	Species
Acanthaceae	<i>Fittonia albivenis</i>
	<i>Justicia pectoralis</i>
Apocynaceae	<i>Tabernaemontana sananho</i>
Aquifoliaceae	<i>Ilex guayusa</i>
Cyperaceae	<i>Cyperus</i> sp.
Fabaceae	<i>Anadenanthera colubrina</i>
	<i>Anadenanthera peregrina</i>
	<i>Calliandra angustifolia</i>
	<i>Pithecellobium unguis-cati</i>
Malpighiaceae	<i>Banisteriopsis caapi</i>
Menispermaceae	<i>Abuta grandifolia</i>
Myristicaceae	<i>Osteophloeum</i> <i>platyspermum</i>
	<i>Virola duckei</i>
	<i>Schuurmansia henningsii</i>
Orchidaceae	<i>Dendrobium pulchellum</i>
	<i>Dendrobium</i> spp.
Rubiaceae	<i>Psychotria carthagenensis</i>
	<i>Psychotria viridis</i>
Solanaceae	<i>Brugmansia</i> × <i>insignis</i>
	<i>Brugmansia suaveolens</i>
	<i>Brugmansia versicolor</i>

Brugmansia sp.

Nicotiana tabacum

4. Discussion

Hunting dogs are indispensable to many traditional cultures. The relationship between these animals and humans is far deeper than in modern cultures. This is evident by both the beliefs regarding dogs and by the ethnoveterinary pharmacopoeias devoted to dogs. Under the influence of the hallucinogenic beverage natem (*Banisteriopsis caapi*), one Shuar shaman sees the commonly-reported vision of boas and jaguars, but also dogs (Bennett et al. 2002). The Quichua believe that dogs have souls and that they dream (Kohn 2007).

The ethnopharmacopoeia for dogs has recently received greater attention from researchers (e.g., Bennett et al. 2002; Lans et al 2000, 2001; Leonard et al. 2002; Jernigan 2009). Yet relatively few studies have examined practices that are said to improve hunting abilities. Koster (2009) summarizes non-pharmacological practices to enhance hunting abilities, which include exposing dogs to the flesh or hair of prey before hunting trips and covering a dog with blood from killed prey, rubbing it with stomach contents, or giving it meat from prey animals. Quichua feed agouti bile or its sternum to their dogs so that they can find this prized game species.

Relatively little attention has been given to plants that are said to improve hunting efficiencies in dogs. Yet their use is widespread. Among the Matsigenka of Peru, hunting ability is believed to be acquired only by the use of plants that enhance a hunter's visual acuity, sense of smell, aim, stamina and luck (Shepard 2002). One quarter of Matsigenka medicinal plant

species are used as hunting medicines and these include plants given to hunting dogs to increase olfactory sensitivity.

The effects of psychoactive substances in humans to other animals is not always comparable. Not surprisingly, nonhuman primates provide are most similar in their responses (Weerts et al. 2007). Dogs respond to commonly-used hallucinogens in a similar manner to humans. Vaupel et al. (1978a) showed that beta-phenethylamine and d-amphetamine increased respiration, dilated pupils and produced restlessness in chronically spinalized dogs. Frith et al. 1987 recorded circling, dilated pupils, hyperactivity, rapid breathing, and salivation in dogs given methylenedioxymethamphetamine. These effects potentially could enhance a dog's hunting ability. Scopolamine significantly impaired memory performance of old, but not young dogs (Araujo et al. 2011). Young hunting dogs seem to be preferred by lowland Amazonian people. For many of the plant compounds, pharmacological studies on dogs is lacking. Nonetheless, based on descriptions of the plants uses, their effects on humans, and their phytochemical profiles, one can speculate on their pharmacological effects in canines.

4.1 Ophthalmics

Only two species were cited as nocturnal ophthalmics – agents that improve vision. Wilbert (1987) reported that a mixture of tobacco and ginger is applied to the eyes of both hunters and their dogs to improve night vision. Few studies have examined the effects of plant extracts on night vision. Tetrahydrocannabinol from *Cannabis sativa* L. has been shown to enhance night visions in some studies (e.g., Russo et al. 2004). The effects of tobacco are less clear. While some studies have shown that tobacco smoke decreases night vision, others have

shown that nicotine enhances night vision, presumably due to the stimulating effects of nicotine (Anonymous 2011). There are no studies on the effects of ginger on night vision. Though the atropine containing genus *Brugmansia* was one of the more frequently cited psychoactive plants given to hunting dogs, the reason for its use was never explicitly said to be related to improvement of vision. Atropine is well-known as a mydriatic and homatropine has been shown to improve nocturnal myopia (Koomen et al. 1951).

4.2 Depuratives/Deodorants

Depuratives/deodorants plants remove or mask odors and improve the ability of hunting dogs to avoid detection by game. Most of them also possess antimicrobial activity that could prevent the development of characteristic odors associated with infections. Shephard (1999) offers an alternate reason for the use of the sulfurous odor of *Mansoa* among the Matsigenka of Peru. They believe that the plant's odor is similar to the smell of peccaries, a preferred Matsigenka game animal.

Other reasons for the use of depuratives/deodorants that transcend the physical effects of moderating odors. The exceedingly fragrant *Siparuna* and *Piper* spp., which are used to bathe dogs, are also employed by the Quichua to cleanse the human body of malevolent spirits. This practice is concordant with their belief that dogs possess souls.

4.3 Olfactory Sensitizers

Little is known about the mechanism of action of Araceae species on olfaction. Most members of the family contain irritating calcium oxalate crystals (Prychid and Rudall 1999).

Another irritant that is employed to enhance olfaction is *Capsicum annuum*. Frasnelli et al. (2009) showed that capsaicins can enhance olfaction in humans and perhaps the same activity is true for canines. Another plant that was said to enhance olfaction was *Tabernaemontana sananho*. Latex from the fruits or leaves are applied to a dog's nose so that "it can smell far." Based on the chemistry of the genus it is likely that, the activity targeting the brain rather than directly affecting the nose. In addition to Jernigan's (2009) study, Brown (1981) reported the use of *Tabernaemontana sananho* for dogs among the Aguaruna and Vickers and Plowman (1984) noted similar use among the Secoya of Ecuador.

4.4. Psychoactives

The use of psychoactive plants given to improve the hunting abilities of dogs is counter-intuitive. Both the Shuar and Quichua give *Ilex guayusa*, which contains methylxanthines, to their hunting dogs. The methylxanthine alkaloid theobromine is toxic to dogs (Strachan and Bennet, 1994). Small doses of the related alkaloid caffeine generate benign arrhythmias in dogs; higher doses cause severe arrhythmias (Mehta et al. 1997). There is clearly a dose-dependent response in canines. Small doses may induce alertness in habituated animals. Quichua deliver small nasal doses to their dogs.

The use of psychoactive hallucinogenic preparations is even more beguiling. The use of hallucinogenic plants in hunting medicines is widespread, but best developed in Ecuador and Peru. All the major Shuar hallucinogens, with the exception of *Brunfelsia grandifolia* also are given to hunting dogs. These hallucinogens contain serotonin agonists (ergot-like alkaloids from *Cyperus* sp.; N-N, dimethyltryptamine from *Virola* and *Psychotria*), monoamine oxidase

inhibitors (β -carboline alkaloids from *Banisteriopsis caapi*), and anticholinergics (scopolamine and atropine from *Brugmansia* spp.) (McKenna et al. 1998, Perry and Perry 1995, Shen et al. 2010).

Serotonin agonists can induce hallucinations in both humans and animals (Wink 2000). Scopolamine hallucinations result from antagonism of ACh receptors (Palfai and Jankiewicz 1997) and the ensuing hallucinations usually often are visual (Perry and Perry, 1995). A common cerebral anticholinergic effect includes bizarre and aggressive behavior (Palfai and Jankiewicz 1997). Acetylcholine plays a key role in sustained attention in both serial reaction time and signal detection tasks. The muscarinic cholinergic antagonist scopolamine substantially impairs accuracy of subjects in the tests (Levin et al. 2011). Aghajanian and Marek (1999) argue that the effects of hallucinogens on glutamatergic transmission in the cerebral cortex may be responsible for higher-level cognitive, perceptual, and affective distortions.

4.4.1 Pharmacology

How then can hallucinogens enhance hunting? Much of the psychopharmacological literature suggests that they are more likely to impair ability. Yet, some evidence is supportive. The mode of action of psilocybin, another serotonin agonist, includes thalamic down-regulation and frontal hypermetabolism, which may contribute to drug-induced synesthesia (Stevenson and Tomiczek, 2007). Synesthesia refers to the stimulation of one sensory or cognitive pathway leads to automatic, involuntary experiences in a second sensory or cognitive pathway. This

process may be responsible not only for hallucinations but also for the enhanced olfactory, auditory or visual senses in hunting dogs.

4.4.2 Case Studies

Other evidence comes from case studies of humans under the influence of hallucinogens. Pedro Kunkumas, a Shuar shaman, said that under the influence of natem (*Banisteriopsis caapi* and *Diplopterys cabrerana* (Cuatrec.) B. Gates) beverages, the body of his patient appears like an x-ray. He can then locate tsentsaks, magical arrows that cause illness, and thus diagnose the patient. Another shaman said that he could hear voices from a long distance under after drinking natem (Bennett et al. 2002). After taking LSD for the first time, two college associates of one of the authors, awoke simultaneously. Both were delirious, believing they were about to be run over by a tractor-trailer. Several hours later they recovered and said that it sounded as if the truck was nearly on top of them. Turning on the lights contributed to their hallucination. Under normal condition, the sound of a trucks downshifting or upshifting on a nearby grade was barely audible. LSD appeared to enhance auditory perception just as natem enhances perceptions in shamans.

Jernigan (2009) reports that the Aguaruna give plants (*Brugmansia* sp., *Mansoa* sp., and *Tabernaemontana sananho*) to their dogs because they produce visions of the intended prey. The Shuar believe that *Brugmansia* spp. help dogs obtain supernatural power (Bennett et al. 2002). For the Quichua described by Kohn (2009), the hallucinogenic mixture tsicta (which includes *Tabernaemontana sananho*, *Nicotiana tabacum*, and *Brugmansia* sp.) is given to dogs so that they can communicate with their masters and to counsel them.

5. Conclusions

Why give psychoactive plants to hunting dogs?

The pharmacological literature suggests equivocal effects of hallucinogens on perception. Moreover, data on the effects of psychoactive plants on dogs is limited deriving mostly from the use of dogs as laboratory surrogates for humans. No studies have investigated the effects of traditional preparations on hunting dogs and the possibility that they can somehow enhance perception that affects the ability to tracking of game animals.

Nonetheless, the practice of administering psychoactive plants to canines is well-established. Could such a practice persist if it impaired hunting success? This is unlikely as hunting is a crucial complement to subsistence practices in the lowland tropics. Vollenweider (1994) hypothesized a disruptive effect of activity of psychedelic substances on sensory gating—the filtering of redundant or superfluous stimuli. Riba et al. (2002), in contrast, suggest ayahuasca has a P50 suppressing effect on sensory gating in humans. We hypothesize that hallucinogenic plants alter perception in hunting dogs by diminishing ancillary signals and enhancing others that aid in the detection and capture of game (Fig. 8). If this is true, the implications are significant. Perhaps plant substance could enhance the ability of dogs to detect explosives, drugs, human remains, or enhance the scores of other abilities for which dogs are valued.




Figure 8. Modified Sensation Model. In normal circumstances, arrows representing sensory input would be of the same magnitude. Under the influence of psychoactive plant extracts, we hypothesize that non-hunting related sensations are blocked, while those related to hunting are enhanced. In tropical rainforests this is most likely to be olfaction.

Acknowledgments

The United States Agency for International Development grants 518-0023-G-SS-4100-00-605 and LAC 0605-G-SS-7037-00 and the Rockefeller Foundation, Program for Economic Botany in Latin America and the Caribbean supported field work for this project. We thank our Shuar and Quichua colleagues for their longstanding collaboration.

References

- Abdullah, Y., Schneider, B., Petersen, M. 2008. Occurrence of rosmarinic acid, chlorogenic acid and rutin in Marantaceae species. *Phytochemistry Letters*. 1, 199-203.
- Aghajanian, G.K., Marek, G.J. 1999. Serotonin and hallucinogens. *Neuropsychopharmacology*. 21, 16S-23S.
- Agbonon, A., Eklugadegboku, K., Aklikokou, K., Gbeassor, M., Akpagana, K., Tam, T.W., Arnason, J.T., Foster, B.C. 2010. In vitro inhibitory effect of West African medicinal and food plants on human cytochrome P450 3A subfamily. *Journal of Ethnopharmacology*. 128, 390-394.
- Ajaiyeoba, E.O., Ekundayo, O. 1999. Essential oil constituents of *Aframomum melegueta* (Roscoe) K. Schum. seeds (alligator pepper) from Nigeria. *Flavour and Fragrance Journal*. 14, 109-111.
- Alarcón, R. 1988. Etnobotánica de los Quichuas de la Amazonia Ecuatoriana. *Miscelánea Antropológica Ecuatoriana*. 7, 1-177.

- Alécio, A.C., da Silva Bolzani, V., Young, M.C., Kato, M.J., Furlan, M. 1998. Antifungal amide from leaves of *Piper hispidum*. *Journal of Natural Products*. 61, 637-639.
- Andrade-Cetto, A., Cardenas Vazquez, R. 2010. Gluconeogenesis inhibition and phytochemical composition of two *Cecropia* species. *Journal of Ethnopharmacology*. 130,93–97.
- Andrade, E.H., Carreira, L.M., da Silva, M.H., da Silva, J.D., Bastos, C.N., Sousa, P.J., Guimarães, E.F., Maia, J.G. 2008. Variability in essential-oil composition of *Piper marginatum* sensu lato. *Chemistry and Biodiversity*. 5, 197-208.
- Angiosperm Phylogeny Group. 2009. An update of the Angiosperm Phylogeny Group classification for the orders and families of flowering plants: APG III. *Botanical journal of the Linnean Society*. 161, 105-121.
- Anonymous, 2009a. *Dendrobiums*. *New Mexico Orchid Guild Newsletter*. 8, 1-3.
- Anonymous. 2009b. Medicinal Plants for Livestock - *Petiveria alliacea*. Cornell University Department of Animal Sciences.
<http://www.ansci.cornell.edu/plants/medicinal/anamu.html> [14 Sep 2014]
- Anonymous, 2011. The Eye and Night Vision - American Optometric Association.
<http://www.aoa.org/optometrists/tools-and-resources/clinical-care-publications/aviation-vision/the-eye-and-night-vision?sso=y> [20 Feb 2015]
- Araujo, J.A., Nobrega, J.N., Raymond, R., Milgram, N.W. 2011. Aged dogs demonstrate both increased sensitivity to scopolamine impairment and decreased muscarinic receptor density. *Pharmacology, Biochemistry, and Behavior*. 98, 203-209.
- Auvin-Guette, C., Baragney, C., Blond, A., Pousset, J.-L., Bodo, B. 1997. Cyclogossine B, a cyclic octapeptide from *Jatropha gossypifolia*. *Journal of Natural Products*. 60, 1155-1157.

- Barbosa-Filho, J.M., Yoshida, M., Gottlieb, O.R. 1989. Lignoids from *Nectandra amazonum* and *N. glabrescens*. *Phytochemistry*. 28, 1991.
- Bennett, B.C. 1992a. Hallucinogenic plants of the Shuar and related indigenous groups in Amazonian Ecuador and Peru. *Brittonia*. 44, 483-493.
- Bennett, B.C. 1992b. Plants and people of the Amazonian rainforests: The role of ethnobotany in sustainable development. *BioScience*. 42, 599-607.
- Bennett, B.C. 1995. Ethnobotany and economic botany of epiphytes, lianas, and other host-dependent plants: An overview, in Lowman, M., Nadkarni, N. (Eds.), *Forest canopies: A review of research on this biological frontier*. Academic Press, New York, pp. 547-586.
- Bennett, B.C. 2007. Doctrine of Signatures: An explanation of medicinal plant discovery or dissemination of knowledge? *Economic Botany*. 61, 246-255.
- Bennett, B.C. 2008. Doctrine of Signatures through Two Millennia. *Herbalgram*. 78, 34-45.
- Bennett, B.C. Baker, M.A., Gómez, P. 2002. Ethnobotany of the Shuar of Eastern Ecuador. *Advances in Economic Botany*. 14, 1-299.
- Bennett, B.C. Alarcón, R. 1994. *Osteophloeum platyspermum* (A.DC.) Warburg and *Virola duckei* A.C. Smith (Myristicaceae): New hallucinogens from Amazonian Ecuador. *Economic Botany*. 48, 152-158.
- Bernart, M.W., Cardellina, J.H., II, Balaschak, M.S., Alexander, M.R., Shoemaker, R.H., Boyd, M.R. 1999. Cytotoxic falcarinol oxylipins from *Dendropanax arboreus*. *Journal of Natural Products*. 59, 748-753.
- Beyer, H. 1908. The symbolic meaning of the dog in ancient Mexico. *American Anthropologist*. 10, 419-422.

- Blumenthal, M., Goldberg, A., Brinckmann, J. (eds.). 2000. Herbal Medicine: Expanded Commission E Monographs. American Botanical Council, Austin, TX.
- Brown, M.F. 1981. Magic and meaning in the world of the Aguaruna Jivaro of Peru. Ph.D. dissertation, University of Michigan, Ann Arbor, Michigan.
- Buskuhl, H., de Oliveira, F.L., Blind, L.Z., de Freitas, R.A., Barison, A., Campos, F.R., Corilo, Y.E., Eberlin, M.N., Caramori, G.F., Biavatti, M.W. 2010. Sesquiterpene lactones from *Vernonia scorpioides* and their in vitro cytotoxicity. *Phytochemistry*. 71, 1539–1544.
- Calixto, J.B., Adair, R., Santos, S., Cechinel Filho, V., Yunes, R.A. 1998. A review of the plants of the genus *Phyllanthus*: Their chemistry, pharmacology, and therapeutic potential. *Medicinal Research Reviews*. 18, 225-258.
- Callaway, J.C., Brito, G.S., Neves, E.S. 2005. Phytochemical analyses of *Banisteriopsis caapi* and *Psychotria viridis*. *Journal of Psychoactive Drugs*. 37, 145-150.
- Carvalho, M.G. de, Yoshida, M., Gottlieb, O.R., Gottlieb, H.E. 1986. Lignans from *Nectandra turbacensis*. *Phytochemistry*. 26, 265-267.
- Chen, C.-H., Chan, H.-Ch., Chu, Y.-T., Ho, H.-Y., Chen, P.-Y., Lee, T.-H., Lee, Ch.-K. 2009. Antioxidant activity of some plant extracts towards xanthine oxidase, lipoxygenase and tyrosinase. *Molecules* 14, 2947-2958.
- Chen, Ch-Y, Cheng, Y.-B., Chen, Sh.-Y., Chien, Ch.-T., Kuo, Y.-H., Guh, J.-H., Khalil, A.T., Shen, Y.-Ch. 2008. New bioactive clerodane diterpenoids from the roots of *Casearia membranacea*. *Chemistry and Biodiversity*. 5, 162-167.
- Cichewicz, R.H., Thorpe, P.A. 1996. The antimicrobial properties of chile peppers (*Capsicum* species) and their uses in Mayan medicine. *Journal of Ethnopharmacology* 52, 61-70.

- Colombo, R., Yariwake, J.H., Queiroz, E.F., Ndjoko, K., Hostettmann, K. 2006. On-line identification of further flavone C- and O-glycosides from sugarcane (*Saccharum officinarum* L., Gramineae) by HPLC-UV-MS. *Phytochemical Analysis*. 17, 337-343.
- Coppinger, R., Schneider, R. 1995. Evolution of working dogs, in Serpell, J. (Ed.), *The domestic Dog: Its Evolution, Behaviour and Interactions with People*. Cambridge University Press, New York. Pp 21-47.
- Costa, P.M., da, Carvalho, M.G., de. 2003. New triterpene isolated from *Eschweilera longipes* (Lecythidaceae). *Anais da Academia Brasileira de Ciências*. 75, 21-25.
- de Diaz, A.M., Gottlieb, O.R. 1979. Propiophenones from *Piper marginatum*. *Planta Medica*. 35, 190-191.
- Diamond, J. 1997. *Guns, Germs, and Steel: The Fates of Human Societies*. W.W. Norton & Co, New York.
- Dong, M.L., Chen, G., Zhou, Z.M. 2008. Flavonoid constituents from *Spiranthes australis* Lindl. *Chemical and Pharmaceutical Bulletin*. 56, 1600-1603.
- Duarte-Almeida, J.M., Novoa, A.V., Linares, A.F., Lajolo, F.M., Inés Genovese, M. 2006. Antioxidant activity of phenolics compounds from sugar cane (*Saccharum officinarum* L.) juice. *Plant Foods for Human Nutrition*. 61, 187-192.
- Duarte-Almeida, J.M., Negri, G., Salatino, A., de Carvalho, J.E., Lajolo, F.M. 2007. Antiproliferative and antioxidant activities of a tricin acylated glycoside from sugarcane (*Saccharum officinarum*) juice. *Phytochemistry* 68, 1165-1171.

- Duke, J.A. 1993. *Jatropha curcas* L. Handbook of Energy Crops.
[http://www.hat.purdue.edu/newcrop/duke-energy/Jatropha curcas.html](http://www.hat.purdue.edu/newcrop/duke-energy/Jatropha%20curcas.html). [accessed 15 June 2014]
- Fatope, M., Takeda, Y., Yamashita, H., Okabe, H., Yamauchi, T. 1990. New cucurbitane triterpenoids from *Momordica charantia*. *Journal of Natural Products*. 53, 1491-1497
- Fernández, A. 1992. Dioses Prehispánicos de México. Panorama Editorial, Mexico City.
- Frith, C.H., Chang, L.W., Lattin, D.L., Walls, R.C., Hamm, J., Doblin, R. 1987. Toxicity of methylenedioxymethamphetamine (MDMA) in the dog and the rat. *Fundamental and Applied Toxicology*. 9, 110-119.
- Flavelle, A. 1991. A traditional agroforestry landscape on Fergusson Island, Papua New Guinea. M.S. Thesis. University of British Columbia. Vancouver, B.C., Canada.
- Frasnelli, J., Oehr, C., Jones-Gotman, M. 2009. Effects of oral irritation on olfaction. *Food Chemistry*. 113, 1003-1007.
- Friedrich, U., Siems, K., Solis, P.N., Gupta, M.P., Jenett-Siems, K. 2005. New prenylated benzoic acid derivatives of *Piper hispidum*. *Die Pharmazie*. 60, 455-457.
- Gadalla, M. 2001. Egyptian Divinities: The All Who are the One. Tehuti Research Foundation, Greensboro, N.C.
- Geron, C., Guenther, A., Greenberg, J., Loescher, H.W., Clark, D., Baker, B. 2002. Biogenic volatile organic compound emissions from a lowland tropical wet forest in Costa Rica. *Atmospheric Environment*. 36, 3793-3802.
- Hansen, L., Boll, P.M. 1986. The polyacetylenic falcarinol as the major allergen in *Schefflera arboricola*. *Phytochemistry* 25, 529-530.

- Hare, B., Brown, M., Williamson, C., Tomasello, M. 2002. The domestication of social cognition in dogs. *Science* 298, 1634-1636.
- Hart, L.A. 1995. Dogs as human companions: A review of the relationship, in Serpell, J. (ed.), *The Domestic Dog: Its Evolution, Behaviour and Interactions with People*. Cambridge University Press, New York. Pp. 161-178.
- Heinrich, M., Chan, J., Wanke, S., Neinhuis, C., Simmonds, M.S.J. 2009. Local uses of *Aristolochia* species and content of nephrotoxic aristolochic acid 1 and 2—A global assessment based on bibliographic sources. *Journal of Ethnopharmacology*. 125, 108-144.
- Holdridge, L.R. 1967. *Life Zone Ecology*, revised edition. Tropical Science Center, San José, Costa Rica.
- Jernigan, K.A. 2009. Barking up the same tree: A comparison of ethnomedicine and canine ethnoveterinary medicine among the Aguaruna. *Journal of Ethnobiology and Ethnomedicine*. 5, 33.
- Kijjoo, A., Giesbrecht, A.M., Akisue, M.K., Gottlieb, O.R., Gottlieb, H.E. 1980. 4-Nerolidylcatechol from *Potomorphe[isic]umbellata*. *Planta Medica*. 39, 85-87.
- Kingston, D.G.I., Li, B.T., Ionescu, F. 2006. Plant anticancer agents III: Isolation of indole and bisindole alkaloids from *Tabernaemontana holstii* roots. *Journal of Pharmaceutical Sciences*. 66, 11351-1138.
- Kohn, E. 2007. How dogs dream: Amazonian natures and the politics of transspecies engagement. *American Ethnologist*. 34, 3-24.

- Koomen, M., Scolnik, R., Tousey, R. 1951. A study of night myopia. *Journal of the Optical Society of America*. 41, 80-83.
- Koster, J. 2009. Hunting dogs in the lowland Neotropics. *Journal of Anthropological Research*. 65, 575-610.
- Kubec R., Musah, R.A. 2001. Cysteine sulfoxide derivatives in *Petiveria alliacea*. *Phytochemistry*. 58, 981-985.
- Kumar, S., Malhotra, R., Kumar, D. 2010. *Euphorbia hirta*: Its chemistry, traditional and medicinal uses, and pharmacological activities. *Pharmacognosy Review*. 4, 58-61.
- Lans C., Harper, T., Georges, K., Bridgewater, E. 2000. Medicinal plants used for dogs in Trinidad and Tobago. *Preventive veterinary medicine*. 45, 201-220.
- Lans, C., Harper, T., George, K., Bridgewater, E. 2001. Medicinal and ethnoveterinary remedies of hunters in Trinidad. *BMC complementary and alternative medicine*. 1, 10.
- Leal, M.B., Elisabetsky, E.J. 1996. Absence of alkaloids in *Psychotria carthagenensis* Jacq. (Rubiaceae). *Journal of Ethnopharmacology*. 54, 37-40.
- Leitão, G.G., Simas, N.K., Soares, S.S.V., de Brito, A.P.P., Claros, B.M.G., Brito3333333, T.B.M., Monache, F.D. 1999. Chemistry and pharmacology of Monimiaceae: A special focus on *Siparuna* and *Mollinedia*. *Journal of Ethnopharmacology*. 65, 87-102.
- Ledón, N., Casacó, A., Rodríguez, V., Cruz, J., González, R., Tolón, Z., Cano, M., Rojas, E. 2003. Anti-Inflammatory and analgesic effects of a mixture of fatty acids isolated and purified from sugar cane wax oil. *Planta Medica*. 69, 367-369.
- Leonard, J.A., Wayne, R.K., Wheeler, J., Valadez, R., Guillén, S., Vilà, C. 2002. Ancient DNA evidence for Old World origin of New World dogs. *Science* 298:1613-1616.

- Levin , E.D., Bushnell, P.J., Rezvani, A.H. 2011. Attention-modulating effects of cognitive enhancers. *Pharmacology, Biochemistry, and Behavior*. 99, 146-154.
- Lewis, W.H., Kennelly, E. J., Bass, G.N., Wedner, H.J., Elvin-Lewis, M.P., Fast, D. 1991. Ritualistic use of the holly *Ilex guayusa* by Amazonian Jivaro Indians. *Journal of Ethnopharmacology*. 33, 25–30.
- Liu, D.-L., N-L Wang, Zhang, X., Yao, and X.-Sh. 2011. Three new triterpenoid saponins from *Ardisia crenata*. *Helvetica Chimica Acta*. 94, 693-702.
- Lognay, G., Marlier, M., Haubruge, E., Trevejo, E., Severin, M. 1989. Study of the lipids from *Renalmia alpinia* (Rott.) Maas. *Grasas y Acietes* 40, 351-355.
- Lognay, G., Marlier, M., Severin, M., Haubruge, E., Gibon, V., Trevejo, E. 1991. On the characterization of some terpenes from *Renalmia alpinia* Rott. (Maas) oleoresin. *Flavour and Fragrance Journal*. 6, 87-91.
- Lopes, S.O., Moreno, P.R., Henriques, A.T. 2000. Growth characteristics and chemical analysis of *Psychotria carthagenensis* cell suspension cultures. *Enzyme and Microbial Technology*. 26, 259-264.
- Macrae, W.D., Towers, G.H.N. 1984. *Justicia pectoralis*: A study of the basis for its use as a hallucinogenic snuff ingredient. *Journal of Ethnopharmacology*. 12, 93-111.
- McKenna, D.J., Callaway, J.C., Grob, C.S. 1998. The scientific investigation of ayahuasca: A review of past and current research. *Heffter Review of Psychedelic Research*. 1, 65-76.
- McKenna, D.J., Luna, L.E., G.N. Towers. 1995. Biodynamic Constituents in Ayahuasca Admixture Plants: An Uninvestigated Folk Pharmacopeia, in Schultes, R.E., von Reis, S.

- (Eds.), *Ethnobotany: Evolution of a Discipline*. Dioscorides Press, Portland OR. Pp. 349-361.
- Mehta, A., Jain, A.C., Mehta, M.C., Billie, M. 1997. Caffeine and cardiac arrhythmias. An experimental study in dogs with review of literature. *Acta Cardiologica*. 52, 273-283.
- Menachery, M.D., Cava, M.P. 1980. Grandirubrine, a new tropoloisoquinoline alkaloid. *Heterocycles*. 14, 943-945.
- Michel, J.L., Chen, Y., Zhang, H., Huang, Y., Kronic, A., Orjala, J., Veliz, M., Soni, K.K., Soejarto, D.D., Caceres, A., Perez, A., Mahady, G.B. 2010. Estrogenic and serotonergic butenolides from the leaves of *Piper hispidum* Swingle (Piperaceae). *Journal of Ethnopharmacology*. 129, 220-226.
- Milliken, W. 1999. Ethnobotany of the Yali of West Papua. <http://rbg-web2.rbge.org.uk/ethnobotany/Yali.pdf>
- Moro, J.C., Fernandes, J.B., Vieira, P.C., Yoshida, M., Gottlieb, O.R., Gottlieb, H.E. 1986. Neolignans from *Nectandra puberula*. *Phytochemistry*. 26, 269-272.
- Nobrega Alves, R.R. da, Barboza, R.R.D., de Medeiros Silva Souto, W., 2010. Chapter 10: Plants Used in Animal Health Care in South and Latin America, in Katerere, D.R., Luseba, D. (Eds.), *Ethnoveterinary Botanical Medicine: Herbal Medicines for Animal Health*. CRC Press Boca Raton FL. Pp. 231- 256.
- Núñez, V., Castro, V., Murillo, R., Ponce-Soto, L.A., Merfort, I., Lomonte, B. 2005. Inhibitory effects of *Piper umbellatum* and *Piper peltatum* extracts towards myotoxic phospholipases A2 from *Bothrops* snake venoms: Isolation of 4-nerolidylcatechol as active principle. *Phytochemistry*. 66, 1017-1025.

- Opitz, S., Schneider, B., 2002. Organ-specific analysis of phenylphenalenone-related compounds in *Xiphidium caeruleum*. *Phytochemistry*. 61, 819-825.
- Oscarsson, K.V., Savage, G.P. 2007. Composition and availability of soluble and insoluble oxalates in raw and cooked taro (*Colocasia esculenta* var. Schott [sic]) leaves. *Food Chemistry*. 101, 559-562.
- Palfai, T., Jankiewicz, H. 1997. *Drugs and human behavior*, second ed. Brown and Benchmark, Madison, WI.
- Pelletier, S.W. 1996. *Alkaloids: Chemical and Biological Perspectives*, Vol. 10. Elsevier, New York.
- Peng, J.Y., Han, X., Xu, L.N., Qi, Y., Xu, Y.W., Wu, Q. 2008. Two new prenylated coumarins from *Spiranthes sinensis* (Pers.) Ames. *Journal of Asian Natural Products Research*. 10, 279-283.
- Perry, E.K., Perry, R.H. 1995. Acetylcholine and hallucinations: Disease-related compared to drug-induced alterations in human consciousness. *Brain and Cognition*. 28, 240-258.
- Pinheiro, M.M., Bessa, S.O., Fingolo, C.E., Kuster, R.M., Matheus, M.E., Menezes, F.S., Fernandes, P.D. 2010. Antinociceptive activity of fractions from *Couroupita guianensis* Aubl. leaves. *Journal of Ethnopharmacology*. 127, 407-413.
- Plowman, T.C., Leuchtmann, A., Blaney, C., Clay, K. 1990. Significance of the fungus *Balansia cyperi* infecting Medicinal Species of *Cyperus* (Cyperaceae) from Amazonia. *Economic Botany*. 44, 452-462.

- Pino Benitez, N., Meléndez León, E.M., Stashenko, E.E. 2009. Essential oil composition from two species of Piperaceae family grown in Colombia. *Journal of Chromatographic Science*. 47, 804-807.
- Pires, A.P., de Oliveira, C.D., Moura, S., Dörr, F.A., Silva, W.A., Yonamine, M. 2009. Gas chromatographic analysis of dimethyltryptamine and beta-carboline alkaloids in ayahuasca, an Amazonian psychoactive plant beverage. *Phytochemical Analysis*. 20, 149-153.
- Prychid, C.J., Rudall, P.J. 1999. Calcium oxalate crystals in monocotyledons: A review of their structure and systematics. *Annals of Botany*. 84, 725-739.
- Ramos, L.S., da Silva, M.L., Luz, A.I.R., Zoghbi, M.G.B., Maia, J.G.S. 1986. Essential oil of *Piper marginatum*. *Journal of Natural Products*. 49, 712-713.
- Ravindranath, N., Venkataiah, B., Ramesh, C., Jayaprakash, P., Das, B. 2003. Jatrophene, a novel macrocyclic bioactive diterpene from *Jatropha gossypifolia*. *Chemical and Pharmaceutical Bulletin*. 51, 870-871.
- Riba, J., Rodríguez-Fornells, A., Barbanoj, M.J. 2002. Effects of ayahuasca on sensory and sensorimotor gating in humans as measured by P50 suppression and prepulse inhibition of the startle reflex, respectively. *Psychopharmacology*. 165, 18-28.
- Rodrigues Silva, D., Baroni, S., Svidzinski, A.E., Bersani-Amado, C.A., Cortez, D.A. 2008. Anti-inflammatory activity of the extract, fractions and amides from the leaves of *Piper ovatum* Vahl (Piperaceae). *Journal of Ethnopharmacology*. 116, 569-573.

- Russo, E.B., Merzouki, A., Mesa, J.M., Frey, K.A., Bach, P.J. 2004. Cannabis improves night vision: A case study of dark adaptometry and scotopic sensitivity in kif smokers of the Rif mountains of northern Morocco. *Journal of Ethnopharmacology*. 93, 99-104.
- Sanz-Biset, J., Campos-de-la-Cruz, J., Epiquién-Rivera, M.A., Cañigueral, S. 2009. A first survey on the medicinal plants of the Chazuta Valley (Peruvian Amazon). *Journal of Ethnopharmacology*. 122, 333–362.
- Savage, G.P., Mårtensson, L., Sedcole, J.R. 2008. Composition of oxalates in baked taro (*Colocasia esculenta* var. Schott) leaves cooked alone or with additions of cows milk or coconut milk. *Journal of Food Composition and Analysis*. 22, 83-86.
- Schultes, R.E., Raffauf, R.F. 1990. *The Healing Forest: Medicinal and toxic plants of the Northwest Amazonia*. Dioscorides Press, Portland, Oregon.
- Schultes, R.E. 1998. Antiquity of the use of New World hallucinogens. *Heffter Review of Psychedelic Research* 1, 1-6.
- Shen, H.W., Jiang, X.L., Winter, J.C., Yu, A.M. 2010. Psychedelic 5-methoxy-N,N-dimethyltryptamine: Metabolism, pharmacokinetics, drug interactions, and pharmacological actions. *Current Drug Metabolism*. 11, 659-666.
- Shepard, G. 1999. *Pharmacognosy and the Senses in Two Amazonian Societies*. Ph.D. Dissertation, University of California, Berkeley, Berkeley, CA.
- Shepard, G. 2002. Primates in Matsigenka subsistence and world view, in Fuentes, A., Wolfe, L.D. (Eds.), *Primates Face to Face: The Conservation Implications of Human-Nonhuman Primate Interconnections*. Cambridge University Press, New York. Pp. 101-136.

- Silva, D.R., Endo, E.H., Filho, B.P., Nakamura, C.V., Svidzinski, T.I., de Souza, A., Young, M.C., Ueda-Nakamura, T., Cortez, D.A. 2009. Chemical composition and antimicrobial properties of *Piper ovatum* Vahl. *Molecules*. 14, 1171-1182.
- Southon, I.W., Buckingham, J. 1989. *Dictionary of Alkaloids*. Chapman and Hall, New York.
- Steele, J.C., Simmonds, M.S., Veitch, N.C., Warhurst, D.C. 1999. Evaluation of the anti-plasmodial activity of bisbenzylisoquinoline alkaloids from *Abuta grandifolia*. *Planta Medica*. 65, 413-416.
- Stevenson, R. J., Tomiczek, C. 2007. Olfactory-induced synesthesias: A review and model. *Psychological Bulletin*. 133, 294-309.
- Strachan, T.R., Bennet, A. 1994. Theobromine poisoning in dogs. *Veterinary Research*. 134, 284.
- Suárez, A.I., Oropeza, M., Vásquez, L., Tillett, S., Compagnone, R.S. 2011. Chemical composition of the essential oil of *Croton gossypifolius* from Venezuela. *Natural Product Communications*. 6, 97-99.
- Tabopda, T.K., Ngoupayo, J., Liu, J., Mitaine-Offer, A.-C., Tanoli, S.A.K., Khan, S.N., Ali, M.S., Ngadjui, B.T., Tsamo, E., Lacaille-Dubois, M.-A., Luu, B. 2008. Bioactive aristolactams from *Piper umbellatum*. *Phytochemistry*. 69, 1726-1731.
- The Plant List. 2015. *The Plant List: A working list of all plant species*. <http://www.theplantlist.org/> (19 Feb 2015).
- Thomas, R, Sah, N.K., Sharma, P.B. 2008. Therapeutic biology of *Jatropha curcas*: A mini review. *Current Pharmaceutical Biotechnology*. 9, 315-324.

- Tillequin, F., Paris, M., Jacquemin, H., Paris, R.R. 1978. [Flavonoids from *Piper marginatum* isolation of a new flavonoid, the marginatoside - Article in French]. *Planta Medica*. 33, 46-52.
- TROPICOS, 2015. Missouri Botanical Garden Tropicos database. [<http://www.tropicos.org/>]
- Vaupel, D.B., Nozaki M., Martin, W.R., Bright, D. 1978. Single dose and cross tolerance studies of beta-phenethylamine, d-amphetamine and LSD in the chronic spinal dog. *European Journal of Pharmacology*. 48, 431-437.
- Vickers, W.T., Plowman, T. 1984. Useful plants of the Siona and Secoya Indians of Eastern Ecuador. *Fieldiana Botany, New Series*. 15, 1-63.
- Vollenweider, F.X. 1994. Evidence for a cortical–subcortical imbalance of sensory information processing during altered states of consciousness using positron emission tomography and [18F]fluorodeoxyglucose, in Pletscher A, Ladewig D (Eds.), 50 years of LSD: Current Status and Perspectives of Hallucinogens. Parthenon, London. Pp 67-86.
- Weerts, E.M., Fantegrossi, W.E., Goodwin, A.K. 2007. The value of nonhuman primates in drug abuse research. *Experimental and Clinical Psychopharmacology*. 15, 309-327.
- Wilbert, J. 1987. Tobacco and shamanism in South America. Yale University Press, New Haven, CT.
- Wink, M. 2000. Interference of Alkaloids with Neuroreceptors and Ion Channels, in Rahman, A. (Ed.), *Studies in Natural Products Chemistry, Volume 21. Bioactive Natural Products (Part B)*. Elsevier, Amsterdam, The Netherlands. Pp. 3-122.

Yang S.W., Zhou, B.N., Wisse, J.H., Evans, R., van der Werff, H., Miller, J.S., Kingston, D.G. 1998.

Three new ellagic acid derivatives from the bark of *Eschweilera coriacea* from the Suriname rainforest. *Journal of Natural Products*. 61, 901-906.

Zhou, B.-N., Baj, N.J., Glass, T.E., Malone, S., Werkhoven, M.C.M., van Troon, F., David, Wisse, J.H., Kingston, D.G.I. 1997. Bioactive labdane diterpenoids from *Renalmia alpinia* collected in the Suriname rainforest. *Journal of Natural Products*. 60, 1287-1293.

Zoghbi, M.G.B., Oliveira, J., Guilhon, G.M.S.P. 2009. The genus *Mansoa* (Bignoniaceae): A source of organosulfur compounds. *Revista Brasileira de Farmacognosia*. 19, 795-804.

Accepted manuscript